

## In-Class Problems Week 4, Mon.

### Problem 1.

The *inverse*  $R^{-1}$  of a binary relation  $R$  from  $A$  to  $B$  is the relation from  $B$  to  $A$  defined by:

$$b R^{-1} a \quad \text{iff} \quad a R b.$$

In other words, you get the diagram for  $R^{-1}$  from  $R$  by “reversing the arrows” in the diagram describing  $R$ . Now many of the relational properties of  $R$  correspond to different properties of  $R^{-1}$ . For example,  $R$  is *total* iff  $R^{-1}$  is a *surjection*.

Fill in the remaining entries in this table:

| $R$ is       | iff $R^{-1}$ is |
|--------------|-----------------|
| total        | a surjection    |
| a function   |                 |
| a surjection |                 |
| an injection |                 |
| a bijection  |                 |

*Hint:* Explain what’s going on in terms of “arrows” from  $A$  to  $B$  in the diagram for  $R$ .

### Arrow Properties

**Definition.** A binary relation,  $R$  is

- is a *function* when it has the [ $\leq 1$  arrow **out**] property.
- is *surjective* when it has the [ $\geq 1$  arrows **in**] property. That is, every point in the right-hand, codomain column has at least one arrow pointing to it.
- is *total* when it has the [ $\geq 1$  arrows **out**] property.
- is *injective* when it has the [ $\leq 1$  arrow **in**] property.
- is *bijection* when it has both the [= 1 arrow **out**] and the [= 1 arrow **in**] property.

### Problem 2.

Assume  $f : A \rightarrow B$  is total function, and  $A$  is finite. Replace the  $\star$  with one of  $\leq, =, \geq$  to produce the *strongest* correct version of the following statements:

- (a)  $|f(A)| \star |B|$ .
- (b) If  $f$  is a surjection, then  $|A| \star |B|$ .
- (c) If  $f$  is a surjection, then  $|f(A)| \star |B|$ .

(d) If  $f$  is an injection, then  $|f(A)| \star |A|$ .

(e) If  $f$  is a bijection, then  $|A| \star |B|$ .

**Problem 3.**

Let  $R : A \rightarrow B$  be a binary relation. Use an arrow counting argument to prove the following generalization of the Mapping Rule 1.

**Lemma.** If  $R$  is a function, and  $X \subseteq A$ , then

$$|X| \geq |R(X)|.$$

**Problem 4.**

Let  $A = \{a_0, a_1, \dots, a_{n-1}\}$  be a set of size  $n$ , and  $B = \{b_0, b_1, \dots, b_{m-1}\}$  a set of size  $m$ . Prove that  $|A \times B| = mn$  by defining a simple bijection from  $A \times B$  to the nonnegative integers from 0 to  $mn - 1$ .

**Problem 5. (a)** Prove that if  $A$  surj  $B$  and  $B$  surj  $C$ , then  $A$  surj  $C$ .

(b) Explain why  $A$  surj  $B$  iff  $B$  inj  $A$ .

(c) Conclude from (a) and (b) that if  $A$  inj  $B$  and  $B$  inj  $C$ , then  $A$  inj  $C$ .

(d) According to the official definition,  $A$  inj  $B$  requires a total injective *relation* ( $\geq 1$  out,  $\leq 1$  in). Explain why  $A$  inj  $B$  iff there is a total injective *function* ( $= 1$  out,  $\leq 1$  in) from  $A$  to  $B$ .